

Paper Type: Original Article

The Second-Order Constant-Coefficient Linear Homogeneous Weak Fuzzy Complex Differential Equations (WFC-DEs) and Initial Value Problems (WFC-IVPs)

Lama Razouk^{1,*} , Suliman Mahmoud¹, Mohamad Ali¹

¹ Department of Mathematics, Latakia University (Tishreen), Latakia, Syria; lamarazouk94@gmail.com; suliman_mmn@yahoo.com; mohamadAli@gmail.com.

Citation:

Received: 25 June 2024 Revised: 19 September 2024 Accepted: 18 November 2024	Razouk, L., Mahmoud, S., & Ali, M. (2025). The second-order constant-coefficient linear homogeneous weak fuzzy complex differential equations (WFC-DEs) and initial value problems (WFC-IVPs). <i>Uncertainty discourse and applications</i> , 2(1), 1-16.
--	--

Abstract

This work aims to introduce the concept of the second-order Weak Fuzzy Complex-Differential Equations (WFC-DEs) for the first time. A special isomorphic transformation function could write a WFC-DE as two related Differential Equations (DEs) of the second order concerning their real variables. We study the second-order constant-coefficient linear homogeneous DE in a Weak Fuzzy Complex (WFC) variable with WFC constant coefficients. Also, we study the simple form of this type of WFC-DE with real constant coefficients. Thus, to find the general solution, we use the characteristic equation of the second-order DE. Also, we get a particular solution for the second-order constant-coefficient linear homogeneous Weak Fuzzy Complex-Initial Value Problem (WFC-IVP). To enhance understanding, we provide illustrative examples for each problem discussed.

Keywords: Weak fuzzy complex numbers, Quadratic weak fuzzy complex equation, Weak fuzzy complex functions, Weak fuzzy complex differential equation, Initial value problem.

1 | Introduction

Throughout history, the expansion of number sets has frequently arisen from the necessity to address particular mathematical challenges. Scholars have shown considerable interest in extending the set of real

 Corresponding Author: lamarazouk94@gmail.com

 <https://doi.org/10.48313/uda.v2i1.50>

 Licensee System Analytics. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0>).

numbers $\mathbb{R}$, with these newly formulated sets being characterized as two-dimensional generalizations of the real number system such as:

Complex numbers: $\{u + vi; u, v \in \mathbb{R}; i^2 = -1\}$, dual numbers: $\{u + vi; u, v \in \mathbb{R}; i^2 = 0\}$, Neutrosophic numbers: $\{u + vi; u, v \in \mathbb{R}; i^2 = i\}$, and split-complex numbers: $\{u + vi; u, v \in \mathbb{R}; i^2 = 1\}$.

Fuzzy numbers emerged as an extension of fuzzy set theory, which Lotfi Zadeh [1], [2] introduced in 1965. Fuzzy numbers represent uncertain numerical values, while traditional real numbers have a precise value. Nowadays, the theory of fuzzy sets has found applications across diverse domains, including artificial intelligence, biology, control systems, decision-making [3], [4], internet [5], information technology [4], electric vehicle [6], economics-industry [7], [8], and portfolio investment [9].

In 2023, the WFC set $F_J = \{u + vJ; u, v \in \mathbb{R}; J^2 = t \in]0, 1[\}$ was defined in [10] as a new generalization of classical real numbers set $\mathbb{R}$ with fuzzy operators $J \notin \mathbb{R}$. Also, they solved linear and quadratic WFC equations [10]. Later, they introduced the WFC vector space [11], the inner products defined over it [12]. Then, Alhasan et al. [13] studied the algebraic properties of WFC matrices and linear systems by applying these matrices.

Furthermore, the linear [14] and the nonlinear [15], [16] Diophantine equation in two WFC variables was studied with the concepts of the WFC and anti-WFC Pythagoras triples and Pythagoras quadruples. However, in [17], main notions were introduced in number theory, like division, ordering, and units in the WFC integers set. Alhasan et al. [18] showed that ‘A-Curves’ display the geometrical characterization of the solutions for some vectorial equations defined by Euclidean norms where they described ‘A-Curves’ as circles and spheres in the two and three-dimensional Euclidean spaces, respectively. Many semi-module isomorphisms are demonstrated between the direct product of WFC numbers with itself and the direct product of classical Euclidean vector spaces multiplied by itself. Also, in [19], those isomorphisms are classified as ‘A-Curves’ and related to the WFC ring.

However, WFC numbers and their primary arithmetic operations were programmed by Python [20]. In addition, a special isomorphism transformation function was created in [21] to help move between the WFC set and the real number set. They defined WFC functions, too. After that, Edduweh et al. [22] presented some types of first-order first-degree separable, exact, and linear Weak Fuzzy Complex-Differential Equations (WFC-DEs). In [23], the novel topological space generated by WFC intervals is based on WFC numbers' partially ordered ring structure. Also, it showed the relationship between the WFC intervals and neutrosophic intervals.

Numerous physical phenomena are represented through functions and their derivatives. The first derivative, for instance, signifies the slope or velocity, while the second derivative conveys curvature or acceleration. DEs play a crucial role in various aspects of contemporary life, finding applications in different fields such as biology, medicine, control systems [24], transportation, meteorology [25], telecommunications, and financial modeling [26].

That leads us to pursue an essential area of inquiry that remains underexplored about the DEs in F_J WFC-DEs. Our paper will introduce the second-order WFC-DEs. In Section 2, we will mention some main concepts about WFC numbers, a special isomorphism transformation function, the quadratic WFC equation, WFC functions, and Weak Fuzzy Complex-Ordinary Differential Equations (WFC-ODEs).

In Section 3, we will define the second-order ordinary DEs, introducing the type of linear homogeneous equation with constant coefficients from F_J . Also, we will take the simple case of them with real constant coefficients. We will find the roots of their characteristic equation and construct their general solution with suitable examples. Then, we will give the notion of Weak Fuzzy Complex-Initial Value Problems (WFC-IVPs) of the related second-order DEs in Section 4 and find their particular solutions.

2 | Preliminaries

This section presents important definitions to explain key elements within our analytical framework [10], [13], [21], [22].

Definition 1. The set of WFC numbers was defined as follows:

$$F_J = \{x_0 + x_1 J; x_0, x_1 \in \mathbb{R}, J^2 = t \in]0, 1[\},$$

where 'J' is the WFC operator ($J \notin \mathbb{R}$).

Definition 2. Let φ be the transformation function from F_J to $\mathbb{R} \times \mathbb{R}$, which we define as follows:

$$\varphi: F_J \mapsto \mathbb{R} \times \mathbb{R},$$

$$\varphi(x_0 + x_1 J) = (x_0 + x_1 (-\sqrt{t}), x_0 + x_1 (+\sqrt{t})) = (x_0 - x_1 \sqrt{t}, x_0 + x_1 \sqrt{t}),$$

where $J^2 = t \in]0, 1[\Rightarrow J = \pm\sqrt{t}$, and $x_0, x_1, y_0, y_1 \in \mathbb{R}$ (This map is an isomorphism).

Definition 3. Let $\varphi: F_J \mapsto \mathbb{R} \times \mathbb{R}$ such that $\varphi(X) = (a, b)$, the inverse function of φ is defined as follows:

$$\varphi^{-1}: \mathbb{R} \times \mathbb{R} \mapsto F_J,$$

$$\varphi^{-1}(a, b) = \frac{1}{2}[a + b] + \frac{1}{2\sqrt{t}}J[b - a].$$

Definition 4. Let $X = x_0 + x_1 J, Y = y_0 + y_1 J \in F_J$, we say that $X \leq Y$, if and only if:

$$\begin{cases} x_0 - x_1 \sqrt{t} \leq y_0 - y_1 \sqrt{t}, \\ x_0 + x_1 \sqrt{t} \leq y_0 + y_1 \sqrt{t}. \end{cases}$$

Definition 5. Let $A = a_0 + a_1 J, B = b_0 + b_1 J \in F_J$, we define the interval $[A, B]$ if and only if $A \leq B$, according to the definition of the partial order relation ($\leq$).

I. If $A \not\leq B$, then $[A, B] = \emptyset$.

II. We can understand $[A, B]$ as follows:

$$[A, B] = \{C = c_0 + c_1 J \in F_J; A \leq C \leq B\}.$$

Theorem 1. Let $aX^2 + bX + c = 0$ be a quadratic WFC equation, where $a = a_0 + a_1 J, b = b_0 + b_1 J$ and $c = c_0 + c_1 J \in F_J$, then it is equivalent to:

$$\begin{cases} A_0 X_0^2 + B_0 X_0 + C_0 = 0, \\ A_1 X_1^2 + B_1 X_1 + C_1 = 0, \end{cases}$$

$$a = \varphi^{-1}(A_0, A_1) = \varphi^{-1}(a_0 - a_1 \sqrt{t}, a_0 + a_1 \sqrt{t}), A_0 = a_0 - a_1 \sqrt{t} \in \mathbb{R}, A_1 = a_0 + a_1 \sqrt{t} \in \mathbb{R},$$

$$B = \varphi^{-1}(B_0, B_1) = \varphi^{-1}(b_0 - b_1 \sqrt{t}, b_0 + b_1 \sqrt{t}), B_0 = b_0 - b_1 \sqrt{t} \in \mathbb{R}, B_1 = b_0 + b_1 \sqrt{t} \in \mathbb{R},$$

$$c = \varphi^{-1}(C_0, C_1) = \varphi^{-1}(c_0 - c_1 \sqrt{t}, c_0 + c_1 \sqrt{t}), C_0 = c_0 - c_1 \sqrt{t} \in \mathbb{R}, C_1 = c_0 + c_1 \sqrt{t} \in \mathbb{R}.$$

Definition 6. Let $f: F_J \mapsto F_J$ be a WFC function in one variable, where

$$\varphi(f(X)) = (f_1(x_0 - x_1 \sqrt{t}), f_2(x_0 + x_1 \sqrt{t})); f_1, f_2: \mathbb{R} \mapsto \mathbb{R},$$

Then we say:

- I. f is continuous on F_J if and only if f_1 and f_2 are continuous on $\mathbb{R}$.
- II. f is differentiable on F_J if and only if f_1 and f_2 are differentiable on $\mathbb{R}$, concerning their variables.
- III. f is integrable on F_J if only f_1 and f_2 are integrable on $\mathbb{R}$.

Definition 7. Let $f: F_J \mapsto F_J$ be a differentiable/integrable function on F_J . We define

- I. $f'(X) = \varphi^{-1}(f'_1(x_0 - x_1 \sqrt{t}), f'_2(x_0 + x_1 \sqrt{t}))$.
- II. $\int f(X).dX = \varphi^{-1}(\int f_1.d(x_0 - x_1 \sqrt{t}), \int f_2.d(x_0 + x_1 \sqrt{t}))$.

Definition 8. Let Y be a dependent variable, and $Y = f(X)$ represents an unknown WFC function where the independent variable $X = x_0 + x_1 J$ is a WFC variable. We define the WFC DE of n -th order (the highest derivative appearing in the DE) as the following:

$$\mathcal{F}(X, Y, Y', Y'', \dots, Y^{(n)}) = 0, \quad (1)$$

where $Y^{(n)} = \frac{d^n Y}{dX^n}$, $Y = y_0 + y_1 J = \varphi^{-1}(y_0 - y_1 \sqrt{t}, y_0 + y_1 \sqrt{t}) = f(X) = \varphi^{-1}(f_1(x_0 - x_1 \sqrt{t}), f_2(x_0 + x_1 \sqrt{t}))$ and $f_1, f_2: \mathbb{R} \mapsto \mathbb{R}$.

Definition 9. A function $Y = f(X) = \varphi^{-1}(f_1(x_0 - x_1 \sqrt{t}), f_2(x_0 + x_1 \sqrt{t}))$ is called the general solution of Eq. (1) on $I \subseteq F_J$ if Y is n -times differentiable WFC function on I and satisfies the equation on I , where $f_1, f_2: \mathbb{R} \mapsto \mathbb{R}$, $I = \varphi^{-1}(I_1 \times I_2) \subseteq F_J$ and $X = \varphi^{-1}(x_0 - x_1 \sqrt{t}, x_0 + x_1 \sqrt{t})$, $x_0 - x_1 \sqrt{t} \in I_1 \subseteq \mathbb{R}$, $x_0 + x_1 \sqrt{t} \in I_2 \subseteq \mathbb{R}$.

We aim to study the second-order DEs that play a pivotal role in modeling systems where the relationship between acceleration and displacement is crucial, such as in mechanics and wave propagation.

3 | Second-Order Ordinary Differential Equations in F_J

We will now propose the definition of the Second-order WFC-DEs. Throughout the discussion, $X = \varphi^{-1}(x_0 - x_1 \sqrt{t}, x_0 + x_1 \sqrt{t}) \in F_J$, $x_0 - x_1 \sqrt{t} \in I_1 \subseteq \mathbb{R}$, $x_0 + x_1 \sqrt{t} \in I_2 \subseteq \mathbb{R}$, $I = \varphi^{-1}(I_1 \times I_2) \subseteq F_J$, $Y_0 = y_0 - y_1 \sqrt{t} \in \mathbb{R}$, $Y_1 = y_0 + y_1 \sqrt{t} \in \mathbb{R}$, $Y = \varphi^{-1}(Y_0, Y_1) \in F_J$.

Definition 10. The WFC-DE of the second order is written as follows:

$$\mathcal{F}(X, Y, Y', Y'') = 0, \quad (2)$$

$$\Leftrightarrow \begin{cases} \mathcal{F}_1(x_0 - x_1 \sqrt{t}, y_0 - y_1 \sqrt{t}, y'_0 - y'_1 \sqrt{t}, y''_0 - y''_1 \sqrt{t}) = 0, \\ \mathcal{F}_2(x_0 + x_1 \sqrt{t}, y_0 + y_1 \sqrt{t}, y'_0 + y'_1 \sqrt{t}, y''_0 + y''_1 \sqrt{t}) = 0, \end{cases} \quad \begin{matrix} (3-1) \\ (3-2) \end{matrix} \quad (3)$$

where $\mathcal{F} = \varphi^{-1}(\mathcal{F}_1, \mathcal{F}_2)$, $y_0 - y_1 \sqrt{t} = f_1(x_0 - x_1 \sqrt{t})$, $y_0 + y_1 \sqrt{t} = f_2(x_0 + x_1 \sqrt{t})$.

$$Y' = \frac{dY}{dX} = f'(X) = \varphi^{-1}(f'_1(x_0 - x_1 \sqrt{t}), f'_2(x_0 + x_1 \sqrt{t})),$$

$$Y'' = \frac{d^2 Y}{dX^2} = f''(X) = \varphi^{-1}(f''_1(x_0 - x_1 \sqrt{t}), f''_2(x_0 + x_1 \sqrt{t})),$$

$$\text{and } \frac{d}{dX} = \varphi^{-1}\left(\frac{d}{d(x_0 - x_1 \sqrt{t})}, \frac{d}{d(x_0 + x_1 \sqrt{t})}\right), \quad \frac{d^2}{dX^2} = \varphi^{-1}\left(\frac{d^2}{d(x_0 - x_1 \sqrt{t})^2}, \frac{d^2}{d(x_0 + x_1 \sqrt{t})^2}\right).$$

Definition 11. Let $Y_0 = y_0 - y_1 \sqrt{t} = f_1(x_0 - x_1 \sqrt{t})$ and $Y_1 = y_0 + y_1 \sqrt{t} = f_2(x_0 + x_1 \sqrt{t})$ are the general solutions to Eq. (3-1) on I_1 and Eq. (3-2) on I_2 , respectively, then

$$Y = f(X) = \varphi^{-1}(Y_0, Y_1),$$

is the general solution of Eq. (2) on $I \subseteq F_J$.

In other words, $Y = f(X)$ is the general solution of *Eq. (2)* on $I \subseteq F_J$ if $f(X)$ is a twice differentiable WFC function on I .

One of the simplest types of second-order ordinary DEs is the linear homogeneous equation with constant coefficients. We will concentrate on this type, which has many applications in science and engineering.

3.1| Second-Order WFC Constant-Coefficient Linear Homogeneous Weak Fuzzy Complex Ordinary Differential Equation

Definition 12. The general form of a second-order linear homogeneous WFC ordinary DE with constant coefficients is written as

$$aY'' + bY' + cY = 0, \quad (4)$$

where $a, b, c, Y, a = a_0 + a_1 J, b = b_0 + b_1 J$ and $c = c_0 + c_1 J \in F_J$.

Remark 1. Using φ , we find that *Eq. (4)* is equivalent to two second-order linear homogeneous ordinary DEs with constant coefficients in R

$$\text{Eq. (4)} \xrightarrow{\varphi} \begin{cases} A_0 Y_0'' + B_0 Y_0' + C_0 Y_0 = 0, & (5-1) \\ A_1 Y_1'' + B_1 Y_1' + C_1 Y_1 = 0, & (5-2) \end{cases} \quad (5)$$

where,

$$a = a_0 + a_1 J = \varphi^{-1}(A_0, A_1), b = b_0 + b_1 J = \varphi^{-1}(B_0, B_1) \text{ and } c = c_0 + c_1 J = \varphi^{-1}(C_0, C_1) \in F_J,$$

$$A_0 = a_0 - a_1 \sqrt{t} \in R, A_1 = a_0 + a_1 \sqrt{t} \in R, \quad (5)$$

$$B_0 = b_0 - b_1 \sqrt{t} \in R, B_1 = b_0 + b_1 \sqrt{t} \in R,$$

$$C_0 = c_0 - c_1 \sqrt{t} \in R, C_1 = c_0 + c_1 \sqrt{t} \in R.$$

We will use the characteristic equation to solve *Eq. (4)*.

Definition 13. The characteristic equation of the constant-coefficient second-order linear homogeneous WFC-DE *Eq. (4)* is defined as the quadratic WFC equation.

$$am^2 + bm + c = 0, \quad a, b, c \in F_J, \quad (6)$$

which holds the following,

It is equivalent to the corresponding characteristic equations

$$\begin{cases} A_0 \lambda^2 + B_0 \lambda + C_0 = 0, & (\text{its discriminant } \Delta_1 > 0), & (7-1) \\ A_1 \mu^2 + B_1 \mu + C_1 = 0, & (\text{its discriminant } \Delta_2 > 0). & (7-2) \end{cases} \quad (7)$$

It has 4 roots when its discriminant $\Delta = b^2 - 4ac$:

$$l_1 = \varphi^{-1}(\lambda_1, \mu_1), l_2 = \varphi^{-1}(\lambda_2, \mu_2), l_3 = \varphi^{-1}(\lambda_1, \mu_2), l_4 = \varphi^{-1}(\lambda_2, \mu_1).$$

Proof:

Depending on *Remark 1* and *Theorem 1*, we know that the characteristic equations of second-order DEs *Eq. (5-1)*, *Eq. (5-2)* have the forms *Eq. (7-1)*, *Eq. (7-2)*, respectively,

$$\begin{cases} A_0 \lambda^2 + B_0 \lambda + C_0 = 0, \\ A_1 \mu^2 + B_1 \mu + C_1 = 0, \end{cases} \quad \varphi^{-1}(A_0 \lambda^2 + B_0 \lambda + C_0, A_1 \mu^2 + B_1 \mu + C_1) = \varphi^{-1}(0, 0)$$

$$\begin{aligned}
&\Rightarrow \varphi^{-1}(A_0\lambda^2, A_1\mu^2) + \varphi^{-1}(B_0\lambda, B_1\mu) + \varphi^{-1}(C_0, C_1) = \varphi^{-1}(0,0) \\
&\Rightarrow \varphi^{-1}(A_0, A_1)\varphi^{-1}(\lambda^2, \mu^2) + \varphi^{-1}(B_0, B_1)\varphi^{-1}(\lambda, \mu) + \varphi^{-1}(C_0, C_1) = \varphi^{-1}(0,0) \\
&\Rightarrow am^2 + bm + c = 0,
\end{aligned}$$

where

$$\begin{aligned}
m &= \varphi^{-1}(\lambda, \mu), a, b, c \in F_j, \\
a &= \varphi^{-1}(A_0, A_1) = \varphi^{-1}(a_0 - a_1\sqrt{t}, a_0 + a_1\sqrt{t}), A_0 = a_0 - a_1\sqrt{t} \in R, A_1 = a_0 + a_1\sqrt{t} \in R, \\
b &= \varphi^{-1}(B_0, B_1) = \varphi^{-1}(b_0 - b_1\sqrt{t}, b_0 + b_1\sqrt{t}), B_0 = b_0 - b_1\sqrt{t} \in R, B_1 = b_0 + b_1\sqrt{t} \in R, \\
c &= \varphi^{-1}(C_0, C_1) = \varphi^{-1}(c_0 - c_1\sqrt{t}, c_0 + c_1\sqrt{t}), C_0 = c_0 - c_1\sqrt{t} \in R, C_1 = c_0 + c_1\sqrt{t} \in R.
\end{aligned}$$

Each characteristic equation has its discriminant,

$$\begin{cases} A_0\lambda^2 + B_0\lambda + C_0 = 0, \\ A_1\mu^2 + B_1\mu + C_1 = 0, \end{cases} \xrightarrow{\text{the discriminants}} \begin{cases} \Delta_1 = B_0^2 - 4A_0C_0, \\ \Delta_2 = B_1^2 - 4A_1C_1. \end{cases}$$

We will study the case when $\Delta_1, \Delta_2 > 0 \Rightarrow$ the corresponding solutions of the characteristic equations *Eq. (7-1)*, *Eq. (7-2)* are real and formed respectively as

$$\begin{cases} \lambda_1 = \frac{-B_0 - \sqrt{\Delta_1}}{2A_0}, \lambda_2 = \frac{-B_0 + \sqrt{\Delta_1}}{2A_0} \in R, \\ \mu_1 = \frac{-B_1 - \sqrt{\Delta_2}}{2A_1}, \mu_2 = \frac{-B_1 + \sqrt{\Delta_2}}{2A_1} \in R, \end{cases}$$

$\Rightarrow$ the solutions of the characteristic equations *Eq. (6)* [10].

$$l_1 = \varphi^{-1}(\lambda_1, \mu_1), l_2 = \varphi^{-1}(\lambda_2, \mu_2), l_3 = \varphi^{-1}(\lambda_1, \mu_2), l_4 = \varphi^{-1}(\lambda_2, \mu_1).$$

Theorem 2. Let $aY'' + bY' + cY = 0$ is a second-order constant-coefficients linear homogeneous DE. And its characteristic equation is $am^2 + bm + c = 0$. Then, the general solution of the WFC-ODE has the form

$$Y = M_1 e^{l_1 X} + M_2 e^{l_2 X} + M_3 e^{l_3 X} + M_4 e^{l_4 X}.$$

Proof:

We found that

$$\begin{aligned}
&aY'' + bY' + c \\
&= 0 \\
&\Leftrightarrow \begin{cases} A_0 Y_0'' + B_0 Y_0' + C_0 Y_0 = 0, \\ A_1 Y_1'' + B_1 Y_1' + C_1 Y_1 = 0, \end{cases} \xrightarrow{\text{the characteristic equations}} \begin{cases} A_0 \lambda^2 + B_0 \lambda + C_0 = 0 \text{ (its discriminant } \Delta_1), \\ A_1 \mu^2 + B_1 \mu + C_1 = 0 \text{ (its discriminant } \Delta_2). \end{cases}
\end{aligned}$$

Since $\Delta_1, \Delta_2 > 0$, the characteristic equations have the real different roots, $\begin{cases} \lambda_1, \lambda_2 \in R, \\ \mu_1, \mu_2 \in R. \end{cases}$

Those roots directly influence the behavior of the solutions of the second-order corresponding Ordinary Differential Equations (ODEs) *Eq. (5-1)* and *Eq. (5-2)* [28],

$$\begin{aligned}
&\xrightarrow{\text{the general solutions}} \begin{cases} Y_0 = H e^{\lambda_1 X_0} + I e^{\lambda_2 X_0} \\ Y_1 = G e^{\mu_1 X_1} + K e^{\mu_2 X_1}, \end{cases} H, I, G, \text{ and } K, \text{ are arbitrary constants.}
\end{aligned}$$

Using φ^{-1} , we get the general solution of WFC-DE *Eq. (4)*,

$$\begin{aligned}
Y &= \varphi^{-1}(Y_0, Y_1) = \varphi^{-1}(\text{He}^{\lambda_1 X_0} + \text{Ie}^{\lambda_2 X_0}, \quad \text{Ge}^{\mu_1 X_1} + \text{Ke}^{\mu_2 X_1}) \\
&= \frac{1}{2}(\text{He}^{\lambda_1 X_0} + \text{Ie}^{\lambda_2 X_0} + \text{Ge}^{\mu_1 X_1} + \text{Ke}^{\mu_2 X_1}) + \frac{1}{2\sqrt{t}}J(\text{Ge}^{\mu_1 X_1} + \text{Ke}^{\mu_2 X_1} - \text{He}^{\lambda_1 X_0} - \text{Ie}^{\lambda_2 X_0}).
\end{aligned}$$

We can write this structure as y_1 or y_2 ,

$$\begin{aligned}
y_1 &= \frac{1}{2}((\text{He}^{\lambda_1 X_0} + \text{Ge}^{\mu_1 X_1}) + (\text{Ie}^{\lambda_2 X_0} + \text{Ke}^{\mu_2 X_1})) + \frac{1}{2\sqrt{t}}J((\text{Ge}^{\mu_1 X_1} - \text{He}^{\lambda_1 X_0}) + \\
&(\text{Ke}^{\mu_2 X_1} - \text{Ie}^{\lambda_2 X_0})) \\
&= \varphi^{-1}(\text{He}^{\lambda_1 X_0}, \text{Ge}^{\mu_1 X_1}) + \varphi^{-1}(\text{Ie}^{\lambda_2 X_0}, \text{Ke}^{\mu_2 X_1}) \\
&= \varphi^{-1}(H, G)\varphi^{-1}(e^{\lambda_1 X_0}, e^{\mu_1 X_1}) + \varphi^{-1}(I, K)\varphi^{-1}(e^{\lambda_2 X_0}, e^{\mu_2 X_1}) \\
&= \varphi^{-1}(H, G) e^{l_1 X} + \varphi^{-1}(I, K) e^{l_2 X},
\end{aligned}$$

or

$$\begin{aligned}
y_2 &= \frac{1}{2}((\text{He}^{\lambda_1 X_0} + \text{Ke}^{\mu_2 X_1}) + (\text{Ie}^{\lambda_2 X_0} + \text{Ge}^{\mu_1 X_1})) + \frac{1}{2\sqrt{t}}J((\text{Ge}^{\mu_1 X_1} - \text{Ie}^{\lambda_2 X_0}) + \\
&(\text{Ke}^{\mu_2 X_1} - \text{He}^{\lambda_1 X_0})) \\
&= \varphi^{-1}(\text{He}^{\lambda_1 X_0}, \text{Ke}^{\mu_2 X_1}) + \varphi^{-1}(\text{Ie}^{\lambda_2 X_0}, \text{Ge}^{\mu_1 X_1}) \\
&= \varphi^{-1}(H, K)\varphi^{-1}(e^{\lambda_1 X_0}, e^{\mu_2 X_1}) + \varphi^{-1}(I, G)\varphi^{-1}(e^{\lambda_2 X_0}, e^{\mu_1 X_1}) \\
&= \varphi^{-1}(H, K) e^{l_3 X} + \varphi^{-1}(I, G) e^{l_4 X}.
\end{aligned}$$

The linearity of equation Eq. (4) means that if we have two solutions y_1 and y_2 , then any linear combination of y_1 and y_2 is also a solution of this equation, i.e.

$$\begin{aligned}
Y(X) &= p_1 y_1(X) + p_2 y_2(X) \\
&= p_1(\varphi^{-1}(H, G) e^{l_1 X} + \varphi^{-1}(I, K) e^{l_2 X}) + p_2(\varphi^{-1}(H, K) e^{l_3 X} + \varphi^{-1}(I, G) e^{l_4 X}) \\
&= M_1 e^{l_1 X} + M_2 e^{l_2 X} + M_3 e^{l_3 X} + M_4 e^{l_4 X},
\end{aligned}$$

is a solution for any choice of constants $p_1, p_2, M_1 = p_1 \varphi^{-1}(H, G), M_2 = p_1 \varphi^{-1}(I, K), M_3 = p_2 \varphi^{-1}(H, K), M_4 = p_2 \varphi^{-1}(I, G)$.

Now, let's verify that the solution $Y(X)$ satisfies Eq. (4).

$$\begin{aligned}
&a(M_1 e^{l_1 X} + M_2 e^{l_2 X} + M_3 e^{l_3 X} + M_4 e^{l_4 X})'' + b(M_1 e^{l_1 X} + M_2 e^{l_2 X} + M_3 e^{l_3 X} + \\
&M_4 e^{l_4 X})' + c(M_1 e^{l_1 X} + M_2 e^{l_2 X} + M_3 e^{l_3 X} + M_4 e^{l_4 X}) \\
&= [a(M_1 e^{l_1 X} + M_2 e^{l_2 X})'' + b(M_1 e^{l_1 X} + M_2 e^{l_2 X})' + c(M_1 e^{l_1 X} + M_2 e^{l_2 X})] \\
&+ [a(M_3 e^{l_3 X} + M_4 e^{l_4 X})'' + b(M_3 e^{l_3 X} + M_4 e^{l_4 X})' + c(M_3 e^{l_3 X} + M_4 e^{l_4 X})] \\
&= p_1[a(\varphi^{-1}(H, G) e^{l_1 X} + \varphi^{-1}(I, K) e^{l_2 X})'' + b(\varphi^{-1}(H, G) e^{l_1 X} + \varphi^{-1}(I, K) e^{l_2 X})' + \\
&c(\varphi^{-1}(H, G) e^{l_1 X} + \varphi^{-1}(I, K) e^{l_2 X})] \\
&+ p_2[a(\varphi^{-1}(H, K) e^{l_3 X} + \varphi^{-1}(I, G) e^{l_4 X})'' + b(\varphi^{-1}(H, K) e^{l_3 X} + \varphi^{-1}(I, G) e^{l_4 X})' + \\
&c(\varphi^{-1}(H, K) e^{l_3 X} + \varphi^{-1}(I, G) e^{l_4 X})] \\
&= p_1[ay_1'' + by_1' + cy_1] + p_2[ay_2'' + by_2' + cy_2]
\end{aligned}$$

$$= p_1(0) + p_2(0) \text{ (because } y_1 \text{ and } y_2 \text{ are solutions of Eq. (4))}$$

$$= 0,$$

where $m = \varphi^{-1}(\lambda, \mu)$, $l_1 = \varphi^{-1}(\lambda_1, \mu_1)$, $l_2 = \varphi^{-1}(\lambda_2, \mu_2)$, $l_3 = \varphi^{-1}(\lambda_1, \mu_2)$, $l_4 = \varphi^{-1}(\lambda_2, \mu_1) \in F_J$.

$$a = a_0 + a_1 J = \varphi^{-1}(A_0, A_1) = \varphi^{-1}(a_0 - a_1 \sqrt{t}, a_0 + a_1 \sqrt{t}), A_0 = a_0 - a_1 \sqrt{t} \in R, A_1 = a_0 + a_1 \sqrt{t} \in R,$$

$$b = b_0 + b_1 J = \varphi^{-1}(B_0, B_1) = \varphi^{-1}(b_0 - b_1 \sqrt{t}, b_0 + b_1 \sqrt{t}), B_0 = b_0 - b_1 \sqrt{t} \in R, B_1 = b_0 + b_1 \sqrt{t} \in R,$$

$$c = c_0 + c_1 J = \varphi^{-1}(C_0, C_1) = \varphi^{-1}(c_0 - c_1 \sqrt{t}, c_0 + c_1 \sqrt{t}), C_0 = c_0 - c_1 \sqrt{t} \in R, C_1 = c_0 + c_1 \sqrt{t} \in R.$$

Example 1. $(3 + 2J)Y'' + JY' - (15 + \frac{45}{2}J)Y = 0$.

Its characteristic equation is

$$(3 + 2J)m^2 + Jm - (15 + \frac{45}{2}J) = 0.$$

The two related DEs of the WFC-DE are [10].

$$\begin{aligned} & \begin{cases} (3 - 2\sqrt{t})Y_0'' - \sqrt{t}Y_0' - \left(15 - \frac{45}{2}\sqrt{t}\right)Y_0 = 0 \\ (3 + 2\sqrt{t})Y_1'' + \sqrt{t}Y_1' - \left(15 + \frac{45}{2}\sqrt{t}\right)Y_1 = 0 \end{cases} \xrightarrow[\text{for } J^2=t=\frac{1}{4}]{=} \begin{cases} 2Y_0'' - \frac{1}{2}Y_0' - \frac{15}{4}Y_0 = 0, \\ 4Y_1'' + \frac{1}{2}Y_1' - \frac{105}{4}Y_1 = 0, \end{cases} \\ & \xrightarrow[\text{the characteristic equations}]{=} \begin{cases} 2\lambda^2 - \frac{1}{2}\lambda - \frac{15}{4} = 0 \text{ (its discriminant } \Delta_1 = \frac{121}{4} > 0) \\ 4\mu^2 + \frac{1}{2}\mu - \frac{105}{4} = 0 \text{ (its discriminant } \Delta_2 = \frac{1681}{4} > 0) \end{cases} \Rightarrow \\ & \begin{cases} \lambda_1 = -\frac{5}{4}, \lambda_2 = \frac{3}{2} \in R, \\ \mu_1 = -\frac{11}{4}, \mu_2 = \frac{5}{2} \in R, \end{cases} \\ & \xrightarrow[\text{the general solutions}]{=} \begin{cases} Y_0 = H e^{-\frac{5}{4}X_0} + I e^{\frac{3}{2}X_0}, \\ Y_1 = G e^{-\frac{11}{4}X_1} + K e^{\frac{5}{2}X_1}, \end{cases} \end{aligned}$$

H, I, G, K are arbitrary constants.

Thus

$$l_1 = \varphi^{-1}(\lambda_1, \mu_1) = \varphi^{-1}\left(-\frac{5}{4}, -\frac{11}{4}\right) = \frac{1}{2}\left(-\frac{5}{4} - \frac{11}{4}\right) + J\left(-\frac{11}{4} + \frac{5}{4}\right) = -2 - \frac{3}{2}J,$$

$$l_2 = \varphi^{-1}(\lambda_2, \mu_2) = \varphi^{-1}\left(\frac{3}{2}, \frac{5}{2}\right) = \frac{1}{2}\left(\frac{3}{2} + \frac{5}{2}\right) + J\left(\frac{5}{2} - \frac{3}{2}\right) = 2 + J,$$

$$l_3 = \varphi^{-1}(\lambda_1, \mu_2) = \varphi^{-1}\left(-\frac{5}{4}, \frac{5}{2}\right) = \frac{1}{2}\left(-\frac{5}{4} + \frac{5}{2}\right) + J\left(\frac{5}{2} + \frac{5}{4}\right) = \frac{5}{8} + \frac{15}{4}J,$$

$$l_4 = \varphi^{-1}(\lambda_2, \mu_1) = \varphi^{-1}\left(\frac{3}{2}, -\frac{11}{4}\right) = \frac{1}{2}\left(\frac{3}{2} - \frac{11}{4}\right) + J\left(-\frac{11}{4} - \frac{3}{2}\right) = -\frac{5}{8} - \frac{17}{4}J.$$

Then, the general solution of WFC-DE

$$Y = M_1 e^{(-2-\frac{3}{2}J)X} + M_2 e^{(2+J)X} + M_3 e^{(\frac{5}{8}+\frac{15}{4}J)X} + M_4 e^{(-\frac{5}{8}-\frac{17}{4}J)X},$$

where $p_1, p_2, M_1 = p_1 \varphi^{-1}(H, G), M_2 = p_1 \varphi^{-1}(I, K), M_3 = p_2 \varphi^{-1}(H, K), M_4 = p_2 \varphi^{-1}(I, G)$ are arbitrary constants.

Now, we will study the simple form of this equation with a real constant.

3.2 | Second-Order Linear Homogeneous WFC-ODE with Real Constant Coefficients

The real constant-coefficient second-order linear homogeneous WFC-DE is adopted as

$$aY'' + bY' + cY = 0, \quad a, b, c \in \mathbb{R}.$$

Its characteristic equation $am^2 + bm + c = 0$, where its discriminant $\Delta = b^2 - 4ac$ and its roots are $l_1, l_2, l_3, l_4 \in \mathbb{F}_J$.

Since the coefficients of the quadratic WFC equation Eq. (6) are real a, b and $c \in \mathbb{R}$, the second-order DEs Eq. (5-1), Eq. (5-2) have the following characteristic equations Eq. (8-1), and Eq. (8-2):

$$\begin{cases} aY_0'' + bY_0' + cY_0 = 0, \\ aY_1'' + bY_1' + cY_1 = 0, \end{cases} \xrightarrow{\text{the characteristic equations}} \begin{cases} a\lambda^2 + b\lambda + c = 0, \\ a\mu^2 + b\mu + c = 0. \end{cases} \quad \begin{matrix} (8-1) \\ (8-2) \end{matrix} \quad (8)$$

Eq. (8-1), and Eq. (8-2) have the discriminants $\Delta_1 = \Delta_2 = \Delta = b^2 - 4ac > 0$.

Then, we will get the real roots of Eq. (8-1), and Eq. (8-2):

$$\lambda_1 = \mu_1 = \frac{-b - \sqrt{\Delta}}{2a}, \quad \lambda_2 = \mu_2 = \frac{-b + \sqrt{\Delta}}{2a},$$

and

$$\lambda_2 = \mu_2 = \frac{-b + \sqrt{\Delta}}{2a}.$$

Example 2.

$$Y'' - 4Y' + 3Y = 0. \quad (9)$$

The characteristic equation of WFC-DE Eq. (9) is

$$m^2 - 4m + 3 = 0. \quad (10)$$

The two related DEs of Eq. (9) are

$$\begin{cases} Y_0'' - 4Y_0' + 3Y_0 = 0, \\ Y_1'' - 4Y_1' + 3Y_1 = 0, \end{cases} \xrightarrow{\text{the characteristic equations}} \begin{cases} \lambda^2 - 4\lambda + 3 = 0 \\ \mu^2 - 4\mu + 3 = 0 \end{cases} \Rightarrow \Delta_1 = \Delta_2 > 0 \Rightarrow$$

$$\begin{cases} \lambda_1 = 3, \lambda_2 = 1 \in \mathbb{R}, \\ \mu_1 = 3, \mu_2 = 1 \in \mathbb{R}, \end{cases}$$

$$\xrightarrow{\text{The general solutions}} \begin{cases} Y_0 = He^{3X_0} + Ie^{X_0}, \\ Y_1 = Ge^{3X_1} + Ke^{X_1}, \end{cases}$$

H, I, G, K are arbitrary constants.

Then the roots of Eq. (10) are

$$\begin{aligned} l_1 &= \varphi^{-1}(\lambda_1, \mu_1) = \varphi^{-1}(3, 3) = \frac{1}{2}(3 + 3) + \frac{1}{2\sqrt{t}}J(3 - 3) = 3, \\ l_2 &= \varphi^{-1}(\lambda_2, \mu_2) = \varphi^{-1}(1, 1) = \frac{1}{2}(1 + 1) + \frac{1}{2\sqrt{t}}J(1 - 1) = 1, \\ l_3 &= \varphi^{-1}(\lambda_1, \mu_2) = \varphi^{-1}(3, 1) = \frac{1}{2}(3 + 1) + \frac{1}{2\sqrt{t}}J(1 - 3) = 2 - \frac{1}{\sqrt{t}}, \\ l_4 &= \varphi^{-1}(\lambda_2, \mu_1) = \varphi^{-1}(1, 3) = \frac{1}{2}(1 + 3) + \frac{1}{2\sqrt{t}}J(3 - 1) = 2 + \frac{1}{\sqrt{t}}. \end{aligned}$$

Then, the general solution of Eq. (9)

$$Y = M_1 e^{3X} + M_2 e^X + M_3 e^{(2-\frac{1}{\sqrt{t}})X} + M_4 e^{(2+\frac{1}{\sqrt{t}})X},$$

where $p_1, p_2, M_1 = p_1 \varphi^{-1}(H, G), M_2 = p_1 \varphi^{-1}(I, K), M_3 = p_2 \varphi^{-1}(H, K), M_4 = p_2 \varphi^{-1}(I, G)$ are arbitrary constants.

We know that arbitrary constants in any general solution of a DE are determined by specifying initial conditions. For a second-order DE in F_J , we should have two WFC initial conditions to determine the two arbitrary constants p_1 and p_2 , which are equivalent to four initial conditions in R .

4 | Second-Order Constant-Coefficient Linear Homogeneous Weak Fuzzy Complex Initial Value Problems

Definition 14. The second-order WFC-Initial Value Problem consists of finding the solution of a WFC-ODE given two initial conditions.

$$\begin{cases} \mathcal{F}(X, Y, Y', Y'') = 0, \\ Y(\alpha) = v, \quad Y'(\alpha) = \omega, \quad (\text{the initial conditions}) \end{cases}, \alpha \leq X \leq \beta, \quad (11)$$

where a fixed WFC value $(\alpha, v), (\alpha, \omega) \in F_J \times F_J$ are given.

Using φ function for Eq. (11), we get

$$\begin{cases} \varphi^{-1}(\mathcal{F}_1(X_0, Y_0, Y_0', Y_0''), \mathcal{F}_2(X_1, Y_1, Y_1', Y_1'')) = 0 \\ \varphi^{-1}(Y_0(\alpha_0), Y_1(\alpha_1)) = \varphi^{-1}(v_0, v_1), \varphi^{-1}(Y_0'(\alpha_0), Y_1'(\alpha_1)) = \varphi^{-1}(\omega_0, \omega_1) \end{cases}$$

$$\Rightarrow \begin{cases} \begin{cases} \mathcal{F}_1(X_0, Y_0, Y_0', Y_0'') = 0, \\ Y_0(\alpha_0) = v_0, Y_0'(\alpha_0) = \omega_0, \end{cases} & \alpha_0 \leq X_0 \leq \beta_0, \quad (12) \Rightarrow \text{its solution is } Y_0(X_0), \\ \begin{cases} \mathcal{F}_2(X_1, Y_1, Y_1', Y_1'') = 0, \\ Y_1(\alpha_1) = v_1, Y_1'(\alpha_1) = \omega_1, \end{cases} & \alpha_1 \leq X_1 \leq \beta_1, \quad (13) \Rightarrow \text{its solution is } Y_1(X_1). \end{cases}$$

Therefore, the particular solution of the second-order WFC-IVP Eq. (11) is a structure of the related solutions of the second-order initial-value problems Eq. (12) and Eq. (13),

$$Y = \varphi^{-1}(Y_0(X_0), Y_1(X_1)),$$

where

$$X = \varphi^{-1}(X_0, X_1) \in [\alpha, \beta] \subseteq F_J, X_0 = x_0 - x_1 \sqrt{t} \in [\alpha_0, \beta_0] \subseteq R, X_1 = x_0 + x_1 \sqrt{t} \in [\alpha_1, \beta_1] \subseteq R,$$

$$Y = \varphi^{-1}(Y_0, Y_1), Y_0 = y_0 - y_1 \sqrt{t}, Y_1 = y_0 + y_1 \sqrt{t},$$

$$\alpha = \varphi^{-1}(\alpha_0, \alpha_1), \beta = \varphi^{-1}(\beta_0, \beta_1), v = \varphi^{-1}(v_0, v_1), \omega = \varphi^{-1}(\omega_0, \omega_1).$$

We get the particular solution when we substitute the initial conditions in the general solution.

Definition 15. The constant-coefficient second-order linear homogeneous WFC initial value problem (WFC-IVP) is formed as

$$\begin{cases} aY'' + bY' + cY = 0, \\ Y(\alpha) = v, Y'(\alpha) = \omega, \end{cases} \quad \alpha \leq X \leq \beta, \quad (12)$$

$$a = \varphi^{-1}(A_0, A_1), \quad A_0 = a_0 - a_1 \sqrt{t}, A_1 = a_0 + a_1 \sqrt{t},$$

$$b = \varphi^{-1}(B_0, B_1), \quad B_0 = b_0 - b_1 \sqrt{t}, B_1 = b_0 + b_1 \sqrt{t},$$

$$c = \varphi^{-1}(C_0, C_1), \quad C_0 = c_0 - c_1 \sqrt{t}, C_1 = c_0 + c_1 \sqrt{t}.$$

Remark 2. Using φ , we find that Eq. (12) is equivalent to two initial value problems of the same type in R ,

$$\text{Eq. (12)} \Rightarrow \begin{cases} \begin{cases} A_0 Y_0'' + B_0 Y_0' + C_0 Y_0 = 0, \\ Y_0(\alpha_0) = v_0, Y_0'(\alpha_0) = \omega_0, \end{cases} & \alpha_0 \leq X_0 \leq \beta_0, \\ \begin{cases} A_1 Y_1'' + B_1 Y_1' + C_1 Y_1 = 0, \\ Y_1(\alpha_1) = v_1, Y_1'(\alpha_1) = \omega_1, \end{cases} & \alpha_1 \leq X_1 \leq \beta_1, \end{cases} \quad (13)$$

where $p_1, p_2, M_1 = p_1 \varphi^{-1}(H, G), M_2 = p_1 \varphi^{-1}(I, K), M_3 = p_2 \varphi^{-1}(H, K), M_4 = p_2 \varphi^{-1}(I, G)$ are determined constants.

Remark 3. Steps to get the particular solution of the second-order WFC-IVP Eq. (12):

1. Solving the second-order initial-value problems IVPs Eq. (13), we get the specific solutions.

$$\begin{cases} Y_0 = \tilde{H}e^{3X_0} + \tilde{I}e^{X_0}, \\ Y_1 = \tilde{G}e^{3X_1} + \tilde{K}e^{X_1}. \end{cases}$$

$\tilde{H}, \tilde{I}, \tilde{G}, \tilde{K}$ are determined constants.

2. Making a structure of the related particular solutions Y_0 and Y_1 of the first step.
3. $Y(X) = p_1(\varphi^{-1}(\tilde{H}, \tilde{G}) e^{1X} + \varphi^{-1}(\tilde{I}, \tilde{K}) e^{12X}) + p_2(\varphi^{-1}(\tilde{H}, \tilde{K}) e^{13X} + \varphi^{-1}(\tilde{I}, \tilde{G}) e^{14X})$.
4. Substituting the initial conditions $Y(\alpha) = v, Y'(\alpha) = \omega$ to have two WFC linear equations.
5. Solving the two WFC linear equations to get values of p_1, p_2 .
6. Substituting p_1 and p_2 in the structure of the second step.

In other words, to find the arbitrary constants $p_1, p_2, M_1 = p_1 \varphi^{-1}(H, G), M_2 = p_1 \varphi^{-1}(I, K), M_3 = p_2 \varphi^{-1}(H, K), M_4 = p_2 \varphi^{-1}(I, G)$, we have to solve the six following equations:

$$Y_0(\alpha_0) = v_0, Y_0'(\alpha_0) = \omega_0 \text{ (which give the values of } H \text{ and } I \text{)}.$$

$$Y_1(\alpha_1) = v_1, Y_1'(\alpha_1) = \omega_1 \text{ (which give the values of } G \text{ and } K \text{)}.$$

Then

$$Y(\alpha) = v, Y'(\alpha) = \omega \text{ (which give the values of } p_1 \text{ and } p_2 \text{)}.$$

Example 3. Second-order linear Homogeneous WFC-IVP with WFC Constant Coefficients.

$$\begin{cases} (3 + 2J) Y'' + J Y' - (15 + \frac{45}{2} J) Y = 0, & 0 \leq X \leq 1 + J. \\ Y(0) = 0, Y'(0) = 1 + J, \end{cases} \quad (14)$$

This Initial Value Problem (IVP) is equivalent to the two following IVPs:

$$\begin{aligned} \text{Eq. (14)} \Rightarrow & \begin{cases} \begin{cases} (3 - 2\sqrt{t}) Y_0'' - \sqrt{t} Y_0' - \left(15 - \frac{45}{2} \sqrt{t}\right) Y_0 = 0, & 0 \leq X_0 \leq 1 - \sqrt{t} \\ Y_0(0) = 0, Y_0'(0) = 1 - \sqrt{t}, \end{cases} \\ \begin{cases} (3 + 2\sqrt{t}) Y_1'' + \sqrt{t} Y_1' - \left(15 + \frac{45}{2} \sqrt{t}\right) Y_1 = 0, & 0 \leq X_1 \leq 1 + \sqrt{t} \\ Y_1(0) = 0, Y_1'(0) = 1 + \sqrt{t}, \end{cases} \end{cases} \\ \xrightarrow[\text{for } J^2 = \frac{1}{4}]{J^2 = \frac{1}{4}} & \begin{cases} \begin{cases} 2Y_0'' - \frac{1}{2} Y_0' - \frac{15}{4} Y_0 = 0, & 0 \leq X_0 \leq \frac{1}{2} \\ Y_0(0) = 0, Y_0'(0) = \frac{1}{2}, \end{cases} \end{cases} \end{aligned} \quad (15)$$

$$\begin{aligned}
 & \xrightarrow[\text{for } J^2=t=\frac{1}{4}]{\text{the general solutions (Example 1)}} \left\{ \begin{aligned} & 4Y_1'' + \frac{1}{2} Y_1' - \frac{105}{4} Y_1 = 0, \quad 0 \leq X_1 \leq \frac{3}{2}, \\ & Y_1(0) = 0, Y_1'(0) = \frac{3}{2}, \end{aligned} \right. \quad (16) \\
 & \xrightarrow[\text{the general solutions (Example 1)}]{\text{the general solutions (Example 1)}} \left\{ \begin{aligned} & Y_0 = He^{-\frac{5}{4}X_0} + Ie^{\frac{3}{2}X_0}, & Y_0' = -\frac{5}{4}He^{-\frac{5}{4}X_0} + \frac{3}{2}Ie^{\frac{3}{2}X_0}, \\ & Y_1 = Ge^{-\frac{11}{4}X_1} + Ke^{\frac{5}{2}X_1}, & Y_1' = -\frac{11}{4}Ge^{-\frac{11}{4}X_1} + \frac{5}{2}Ke^{\frac{5}{2}X_1}. \end{aligned} \right.
 \end{aligned}$$

After substituting the corresponding initial conditions to find the values of H, I, G, K:

$$\begin{aligned}
 & \left\{ \begin{aligned} & 0 = H + I, \\ & \frac{1}{2} = -\frac{5}{4}H + \frac{3}{2}I, \end{aligned} \right. \xrightarrow[\text{solving this system}]{} \left\{ \begin{aligned} & \tilde{H} = \frac{-2}{11}, \\ & \tilde{I} = \frac{2}{11} \end{aligned} \right. \xrightarrow[\text{the particular solution for IVP Eq. (15)}]{\text{the particular solution for IVP Eq. (15)}} Y_0 = \frac{-2}{11}e^{-\frac{5}{4}X_0} + \frac{2}{11}e^{\frac{3}{2}X_0}, \\
 & \left\{ \begin{aligned} & 0 = G + K, \\ & \frac{3}{2} = -\frac{11}{4}G + \frac{5}{2}K, \end{aligned} \right. \xrightarrow[\text{solving this system}]{} \left\{ \begin{aligned} & \tilde{G} = \frac{-6}{21}, \\ & \tilde{K} = \frac{6}{21} \end{aligned} \right. \xrightarrow[\text{the particular solution for IVP Eq. (16)}]{\text{the particular solution for IVP Eq. (16)}} Y_1 = \frac{-6}{21}e^{-\frac{11}{4}X_1} + \frac{6}{21}e^{\frac{5}{2}X_1}, \\
 & l_1 = \varphi^{-1}(\lambda_1, \mu_1) = \varphi^{-1}\left(-\frac{5}{4}, -\frac{11}{4}\right) = \frac{1}{2}\left(-\frac{5}{4} - \frac{11}{4}\right) + J\left(-\frac{11}{4} + \frac{5}{4}\right) = -2 - \frac{3}{2}J, \\
 & l_2 = \varphi^{-1}(\lambda_2, \mu_2) = \varphi^{-1}\left(\frac{3}{2}, \frac{5}{2}\right) = \frac{1}{2}\left(\frac{3}{2} + \frac{5}{2}\right) + J\left(\frac{5}{2} - \frac{3}{2}\right) = 2 + J, \\
 & l_3 = \varphi^{-1}(\lambda_1, \mu_2) = \varphi^{-1}\left(-\frac{5}{4}, \frac{5}{2}\right) = \frac{1}{2}\left(-\frac{5}{4} + \frac{5}{2}\right) + J\left(\frac{5}{2} + \frac{5}{4}\right) = \frac{5}{8} + \frac{15}{4}J, \\
 & l_4 = \varphi^{-1}(\lambda_2, \mu_1) = \varphi^{-1}\left(\frac{3}{2}, -\frac{11}{4}\right) = \frac{1}{2}\left(\frac{3}{2} - \frac{11}{4}\right) + J\left(-\frac{11}{4} - \frac{3}{2}\right) = -\frac{5}{8} - \frac{17}{4}J, \\
 & \varphi^{-1}\left(\frac{-2}{11}, \frac{-6}{21}\right) = \frac{1}{2}\left(\frac{-2}{11} - \frac{6}{21}\right) + J\left(\frac{-6}{21} - \frac{-2}{11}\right) = -\frac{18}{77} - \frac{8}{77}J, \\
 & \varphi^{-1}\left(\frac{2}{11}, \frac{6}{21}\right) = \frac{1}{2}\left(\frac{2}{11} + \frac{6}{21}\right) + J\left(\frac{6}{21} - \frac{2}{11}\right) = \frac{18}{77} + \frac{8}{77}J, \\
 & \varphi^{-1}\left(\frac{-2}{11}, \frac{6}{21}\right) = \frac{1}{2}\left(\frac{-2}{11} + \frac{6}{21}\right) + J\left(\frac{6}{21} + \frac{2}{11}\right) = \frac{4}{77} + \frac{36}{77}J, \\
 & \varphi^{-1}\left(\frac{2}{11}, \frac{-6}{21}\right) = \frac{1}{2}\left(\frac{2}{11} + \frac{-6}{21}\right) + J\left(\frac{-6}{21} - \frac{2}{11}\right) = -\frac{4}{77} - \frac{36}{77}J.
 \end{aligned}$$

Then, the general solution of Eq. (14) (from Example 1)

$$\begin{aligned}
 Y(X) &= M_1 e^{(-2-\frac{3}{2})X} + M_2 e^{(2+J)X} + M_3 e^{(\frac{5}{8}+\frac{15}{4}J)X} + M_4 e^{(-\frac{5}{8}-\frac{17}{4}J)X} \\
 &= p_1(\varphi^{-1}(\tilde{H}, \tilde{G}) e^{l_1 X} + \varphi^{-1}(\tilde{I}, \tilde{K}) e^{l_2 X}) + p_2(\varphi^{-1}(\tilde{H}, \tilde{K}) e^{l_3 X} + \varphi^{-1}(\tilde{I}, \tilde{G}) e^{l_4 X}) \\
 &= p_1(\varphi^{-1}\left(\frac{-2}{11}, \frac{-6}{21}\right) e^{l_1 X} + \varphi^{-1}\left(\frac{2}{11}, \frac{6}{21}\right) e^{l_2 X}) + p_2(\varphi^{-1}\left(\frac{-2}{11}, \frac{6}{21}\right) e^{l_3 X} + \\
 &\quad \varphi^{-1}\left(\frac{2}{11}, \frac{-6}{21}\right) e^{l_4 X}) \\
 &= p_1\left(\left(-\frac{18}{77} - \frac{8}{77}J\right) e^{(-2-\frac{3}{2})X} + \left(\frac{18}{77} + \frac{8}{77}J\right) e^{(2+J)X}\right) + p_2\left(\left(\frac{4}{77} + \frac{36}{77}J\right) e^{(\frac{5}{8}+\frac{15}{4}J)X} + \right. \\
 &\quad \left. \left(-\frac{4}{77} - \frac{36}{77}J\right) e^{(-\frac{5}{8}-\frac{17}{4}J)X}\right),
 \end{aligned}$$

$$Y'(X) = p_1 \left(\left(-2 - \frac{3}{2}J \right) \left(-\frac{18}{77} - \frac{8}{77}J \right) e^{\left(-2 - \frac{3}{2}J \right)X} + (2+J) \left(\frac{18}{77} + \frac{8}{77}J \right) e^{(2+J)X} \right) +$$

$$p_2 \left(\left(\frac{5}{8} + \frac{15}{4}J \right) \left(\frac{4}{77} + \frac{36}{77}J \right) e^{\left(\frac{5}{8} + \frac{15}{4}J \right)X} + \left(-\frac{5}{8} - \frac{17}{4}J \right) \left(-\frac{4}{77} - \frac{36}{77}J \right) e^{\left(-\frac{5}{8} - \frac{17}{4}J \right)X} \right).$$

Substituting the initial conditions: $Y(0) = 0$ in the general solution Y , and $Y'(0) = 1 + J$ in the first derivative of the general solution Y' , we get

$$\begin{cases} 0 = p_1 \left(\left(-\frac{18}{77} - \frac{8}{77}J \right) + \left(\frac{18}{77} + \frac{8}{77}J \right) \right) + p_2 \left(\left(\frac{4}{77} + \frac{36}{77}J \right) + \left(-\frac{4}{77} - \frac{36}{77}J \right) \right), \\ 1+J = p_1 \left(\left(2 + \frac{3}{2}J \right) \left(\frac{18}{77} + \frac{8}{77}J \right) + (2+J) \left(\frac{18}{77} + \frac{8}{77}J \right) \right) + p_2 \left(\left(\frac{5}{8} + \frac{15}{4}J \right) \left(\frac{4}{77} + \frac{36}{77}J \right) + \left(\frac{5}{8} + \frac{17}{4}J \right) \left(\frac{4}{77} + \frac{36}{77}J \right) \right) \end{cases}$$

$$\Rightarrow \begin{cases} 0 = 0p_1 + 0p_2 \\ 1+J = \left(\frac{18}{77} + \frac{8}{77}J \right) \left(4 + \frac{5}{2}J \right) p_1 + \left(\frac{4}{77} + \frac{36}{77}J \right) \left(\frac{10}{8} + \frac{32}{4}J \right) p_2. \end{cases}$$

We find that those two equations are satisfied when

$$p_1 = \frac{\left[1+J - \left(\frac{4}{77} + \frac{36}{77}J \right) \left(\frac{10}{8} + \frac{32}{4}J \right) p_2 \right]}{\left[\left(\frac{18}{77} + \frac{8}{77}J \right) \left(4 + \frac{5}{2}J \right) \right]} = P(p_2).$$

Then the solution of Eq. (15) is in terms of p_2 for any p_1 :

$$Y(X) = P(p_2) \left(\left(-\frac{18}{77} - \frac{8}{77}J \right) e^{\left(-2 - \frac{3}{2}J \right)X} + \left(\frac{18}{77} + \frac{8}{77}J \right) e^{(2+J)X} \right) + p_2 \left(\left(\frac{4}{77} + \frac{36}{77}J \right) e^{\left(\frac{5}{8} + \frac{15}{4}J \right)X} + \left(-\frac{4}{77} - \frac{36}{77}J \right) e^{\left(-\frac{5}{8} - \frac{17}{4}J \right)X} \right).$$

Now, we will solve a simple form, 'the real constant-coefficient second-order linear homogeneous WFC ordinary DE.'

Example 4. Second-order linear Homogeneous WFC-IVP with Real Constant Coefficients.

$$\begin{cases} Y'' - 4Y' + 3Y = 0, \\ Y(0) = 3, Y'(0) = 5, \end{cases} \quad 0 \leq X \leq 10. \quad (17)$$

This IVP is equivalent to the two following IVPs:

$$\text{Eq. (17)} \Rightarrow \begin{cases} \begin{cases} Y_0'' - 4Y_0' + 3Y_0 = 0, \\ Y_0(0) = 3, Y_0'(0) = 5, \end{cases} & 0 \leq X_0 \leq 10, \\ \begin{cases} Y_1'' - 4Y_1' + 3Y_1 = 0, \\ Y_1(0) = 3, Y_1'(0) = 5, \end{cases} & 0 \leq X_1 \leq 10, \end{cases}$$

$$\xrightarrow[\text{from Example 2}]{\text{the general solutions}} \begin{cases} Y_0 = He^{3X_0} + Ie^{X_0}, Y_0' = 3He^{3X_0} + Ie^{X_0}, \\ Y_1 = Ge^{3X_1} + Ke^{X_1}, Y_1' = 3Ge^{3X_1} + Ke^{X_1}. \end{cases}$$

After substituting the corresponding initial conditions,

$$\begin{cases} \begin{cases} 3 = H + I \\ 5 = 3H + I \end{cases} \Rightarrow \begin{cases} \tilde{H} = 1 \\ \tilde{I} = 2 \end{cases} \Rightarrow Y_0 = e^{3X_0} + 2e^{X_0}, \\ \begin{cases} 3 = G + K \\ 5 = 3G + K \end{cases} \Rightarrow \begin{cases} \tilde{G} = 1 \\ \tilde{K} = 2 \end{cases} \Rightarrow Y_1 = e^{3X_1} + 2e^{X_1}. \end{cases}$$

The general solution of Eq. (17),

$$\begin{aligned} Y(X) &= p_1(\varphi^{-1}(\tilde{H}, \tilde{G}) e^{l_1 X} + \varphi^{-1}(\tilde{I}, \tilde{K}) e^{l_2 X}) + p_2(\varphi^{-1}(\tilde{H}, \tilde{K}) e^{l_3 X} + \varphi^{-1}(\tilde{I}, \tilde{G}) e^{l_4 X}) \\ &= p_1(\varphi^{-1}(1, 1) e^{l_1 X} + \varphi^{-1}(2, 2) e^{l_2 X}) + p_2(\varphi^{-1}(1, 2) e^{l_3 X} + \varphi^{-1}(2, 1) e^{l_4 X}) \end{aligned}$$

$$\varphi^{-1}(1, 1) = \frac{1}{2}(1 + 1) + \frac{1}{2\sqrt{t}}J(1 - 1) = 1,$$

$$\varphi^{-1}(2, 2) = 2,$$

$$\varphi^{-1}(1, 2) = \frac{1}{2}(1 + 2) + \frac{1}{2\sqrt{t}}J(2 - 1) = \frac{3}{2} + \frac{1}{2\sqrt{t}}J,$$

$$\varphi^{-1}(2, 1) = \frac{1}{2}(2 + 1) + \frac{1}{2\sqrt{t}}J(1 - 2) = \frac{3}{2} - \frac{1}{2\sqrt{t}}J,$$

$$l_1 = \varphi^{-1}(\lambda_1, \mu_1) = 3,$$

$$l_2 = \varphi^{-1}(\lambda_2, \mu_2) = 1,$$

$$l_3 = \varphi^{-1}(\lambda_1, \mu_2) = 2 - \frac{1}{\sqrt{t}}J,$$

$$l_4 = \varphi^{-1}(\lambda_2, \mu_1) = 2 + \frac{1}{\sqrt{t}}J.$$

Then

$$Y(X) = p_1(e^{3X} + 2e^X) + p_2\left[\left(\frac{3}{2} + \frac{1}{2\sqrt{t}}J\right)e^{\left(2 - \frac{1}{\sqrt{t}}J\right)X} + \left(\frac{3}{2} - \frac{1}{2\sqrt{t}}J\right)e^{\left(2 + \frac{1}{\sqrt{t}}J\right)X}\right],$$

$$\begin{aligned} Y'(X) &= p_1(3e^{3X} + 2e^X) + p_2\left[\left(2 - \frac{1}{\sqrt{t}}J\right)\left(\frac{3}{2} + \frac{1}{2\sqrt{t}}J\right)e^{\left(2 - \frac{1}{\sqrt{t}}J\right)X} + \left(2 + \frac{1}{\sqrt{t}}J\right)\left(\frac{3}{2} - \frac{1}{2\sqrt{t}}J\right)e^{\left(2 + \frac{1}{\sqrt{t}}J\right)X}\right]. \end{aligned}$$

Substituting the initial conditions: $Y(0) = 1$ in the general solution Y , and $Y'(0) = 1$ in the first derivative of the general solution Y' , we get

$$\begin{aligned} &\begin{cases} 3 = p_1(1 + 2) + p_2\left(\frac{3}{2} + \frac{1}{2\sqrt{t}}J + \frac{3}{2} - \frac{1}{2\sqrt{t}}J\right), \\ 5 = p_1(3 + 2) + p_2\left[\left(2 - \frac{1}{\sqrt{t}}J\right)\left(\frac{3}{2} + \frac{1}{2\sqrt{t}}J\right) + \left(2 + \frac{1}{\sqrt{t}}J\right)\left(\frac{3}{2} - \frac{1}{2\sqrt{t}}J\right)\right], \end{cases} \\ &\Rightarrow \begin{cases} 3 = 3p_1 + 3p_2 \\ 5 = 5p_1 + p_2\left[3 - \frac{1}{\sqrt{t}}J - \frac{3}{2\sqrt{t}}J - \frac{1}{2} + 3 - \frac{1}{\sqrt{t}}J + \frac{3}{2\sqrt{t}}J - \frac{1}{2}\right] \end{cases} \\ &\quad \Rightarrow \begin{cases} 1 = p_1 + p_2 \\ 5 = 5p_1 + \left(5 - \frac{2}{\sqrt{t}}J\right)p_2. \end{cases} \end{aligned}$$

Solving these two equations [12], we find $p_1 = 1, p_2 = 0$ and the particular solution of Eq. (17) is

$$Y(X) = e^{3X} + 2e^X.$$

5 | Conclusion

In this paper, we have defined the second-order WFC-DEs. We have focused on solving the second-order constant-coefficient linear homogeneous WFC DE using its characteristic equation, where we have found its

general solution. We have solved the related second-order constant-coefficient linear homogeneous IVPs by adding initial conditions. We have explained all examples for this type of DEs and IVPs with WFC constant coefficients and real constant coefficients.

Author Contribution

Conceptualization and Methodology, L. R., writing-reviewing and editing L. R., S. M. and M. A. All authors have read and agreed to the published version of the manuscript.

Funding

The authors declare that no external funding or support was received for the research.

Data Availability

All data supporting the reported findings in this research paper are provided within the manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

References

- [1] Zadeh, L. A. (1965). Fuzzy sets. *Information and control*, 8(3), 338–353. [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X)
- [2] Wang-Jin, L. (1983). Operations on fuzzy ideals. *Fuzzy sets and systems*, 11(1), 31–39. <https://dl.acm.org/doi/abs/10.5555/2846983.2847100>
- [3] Sorourkhah, A., & Edalatpanah, S. A. (2022). Using a combination of matrix approach to robustness analysis (MARA) and fuzzy DEMATEL-based ANP (FDANP) to choose the best decision. *International journal of mathematical, engineering and management sciences*, 7(1), 68–80. <https://doi.org/10.33889/IJMEMS.2022.7.1.005>
- [4] Ihsan, M., Saeed, M., & Rahman, A. U. (2023). Optimizing hard disk selection via a fuzzy parameterized single-valued neutrosophic soft set approach. *Journal of operational and strategic analytics*, 1(2), 62–69. <https://doi.org/10.56578/josa010203>
- [5] Abdi, G. H., Refahi Sheikhani, A. H., Kordrostami, S., & Babaie, S. (2023). A novel selfish node detection based on fuzzy system and game theory in internet of things. *Fuzzy optimization and modeling journal*, 4(4), 32–47. <https://doi.org/10.30495/fomj.2024.1996589.1122>
- [6] Ghobadi, A., Tavakkoli Moghadam, R., Fallah, M., & Kazemipoor, H. (2021). Multi-depot electric vehicle routing problem with fuzzy time windows and pickup/delivery constraints. *Journal of applied research on industrial engineering*, 8(1), 1–18. <https://doi.org/10.22105/jarie.2021.231764.1165>
- [7] Rahimi Ghazikalayeh, A., Amirafshari, M., Mkrchyan, H. M., & Taji, M. (2013). Application of fuzzy hybrid analytic network process in equipment selection of open-pit metal mines. *International journal of research in industrial engineering*, 2(3), 35–46. http://www.riejournal.com/article_47953.html
- [8] Halder, P. K., Karmaker, C. L., Kundu, B., & Daniel, T. (2018). Evaluation of factors affecting the productivity of RMG in Bangladesh : A fuzzy AHP approach. *International journal of research in industrial engineering*, 7(1), 51–60. <https://doi.org/10.22105/riej.2017.94987.1011>
- [9] El-Morsy, S. (2023). Stock portfolio optimization using pythagorean fuzzy numbers. *Journal of operational and strategic analytics*, 1(1), 8–13. <https://doi.org/10.56578/josa010205josa010205>
- [10] Hatip, A. (2023). An introduction to weak fuzzy complex numbers. *Galoitica: journal of mathematical structures and applications*, 3(2), 08–13. <https://doi.org/10.54216/gjmsa.030101>
- [11] Hatip, A. (2023). On the fuzzy weak complex vector spaces. *Galoitica: journal of mathematical structures and applications*, 3(1), 29–32. <https://doi.org/10.54216/gjmsa.030105>

- [12] Ali, R. (2023). On the weak fuzzy complex inner products on weak fuzzy complex vector spaces. *Neoma journal of mathematics and computer science*, 1. <https://doi.org/10.5281/zenodo.7953682>
- [13] Alhasan, Y. A., Alfahal, A. M. A., Abdulfatah, R. A., Nordo, G., & Zahra, M. M. A. (2023). On some novel results about weak fuzzy complex matrices. *International journal of neutrosophic science*, 21(1), 134–140. <https://doi.org/10.54216/IJNS.210112>
- [14] Abualhomos, M., Salameh, W. M. M., Bataineh, M., Al-Qadri, M. O., Alahmade, A., & Al-Husban, A. (2024). An effective algorithm for solving weak fuzzy complex diophantine equations in two variables. *International journal of neutrosophic science*, 23(4), 386–394. <https://doi.org/10.54216/IJNS.230431>
- [15] Alfahal, A. M. A., Abobala, M., Alhasan, Y. A., & Abdulfatah, R. A. (2023). Generating weak fuzzy complex and anti weak fuzzy complex integer solutions for pythagoras diophantine equation $X^2 + Y^2 = Z^2$. *International journal of neutrosophic science*, 22(2), 8–14. <https://doi.org/10.54216/IJNS.220201>
- [16] Galarza, F. C., Flores, M. L., Rivero, D. P., & Abobala, M. (2023). On weak fuzzy complex pythagoras quadruples. *International journal of neutrosophic science*, 22(2), 108–113. <https://doi.org/10.54216/IJNS.220209>
- [17] Abobala, M. (2024). On some novel results about weak fuzzy complex integers. *Journal of neutrosophic and fuzzy systems*, 15–22. <https://doi.org/10.54216/JNFS.080202>
- [18] Alhasan, Y. A., Xu, L., Abdulfatah, R. A., & Alfahal, A. M. A. (2023). The geometrical characterization for the solutions of a vectorial equation by using weak fuzzy complex numbers and other generalizations of real numbers. *International journal of neutrosophic science*, 21(4), 155–159. <https://doi.org/10.54216/IJNS.210415>
- [19] Shihadeh, A., Salameh, W. M. M., Bataineh, M., Al-Tarawneh, H., Alahmade, A., & Al-Husban, A. (2024). On the geometry of weak fuzzy complex numbers and applications to the classification of some a-curves. *International journal of neutrosophic science*, 23(4), 369–375. <https://doi.org/10.54216/IJNS.230428>
- [20] Razouk, L., Mahmoud, S., & Ali, M. (2023). A computer program for the system of weak fuzzy complex numbers and their arithmetic operations using Python. *Galoitica: journal of mathematical structures and applications*, 8(1), 45–51. <https://doi.org/10.54216/gjmsa.080104>
- [21] Razouk, L., Mahmoud, S., & Ali, M. (2024). On the foundations of weak fuzzy complex-real functions. *Journal of fuzzy extension and applications*, 5(1), 116–140. <https://doi.org/10.22105/jfea.2024.435955.1369>
- [22] Edduweh, H., Heilat, A. S., Razouk, L., Khalil, S. A., Alsaraireh, A. A., & Al-Husban, A. (2025). On the weak fuzzy complex differential equations and some types of the 1st order 1st degree WFC-ODEs. *International journal of neutrosophic science*, 25(3), 450–468. <https://doi.org/10.54216/IJNS.250338>
- [23] Hatamleh, R. (2025). On a novel topological space based on partially ordered ring of weak fuzzy complex numbers and its relation with the partially ordered neutrosophic ring of real numbers. *Neutrosophic sets and systems*, 78(1), 577–590. https://digitalrepository.unm.edu/nss_journal/vol78/iss1/31/
- [24] Khalil, H. K., & Grizzle, J. W. (2002). *Nonlinear systems*. Prentice Hall. <https://testbankdeal.com/sample/nonlinear-systems-3rd-edition-khalil-solutions-manual.pdf>
- [25] Warner, T. T. (2010). *Numerical weather and climate prediction*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511763243>
- [26] John C Hull, S. B. (1988). *Options, futures, and other derivatives*. Pearson India. [file:///C:/Users/NoteBook/Downloads/Options,%20Futures%20and%20Other%20Derivatives%208th%20John%20Hull%20\(1\).pdf](file:///C:/Users/NoteBook/Downloads/Options,%20Futures%20and%20Other%20Derivatives%208th%20John%20Hull%20(1).pdf)
- [27] Burden, R. L. & Faires, J. D. (2010). *Numerical analysis*. Cengage Learning. <https://pdfcoffee.com/numerical-analysis-9th-edition-pdf-free.html>
- [28] Egede, U. (2018). *Complex analysis & differential equations*. <https://marksphysicshelp.com/wp-content/uploads/2019/04/complexanalysisnotes2.pdf>